

FAQ Coronavirus (COVID-19)

Updated 21 January 2022

Table of Contents

Updated 21 January 2022	1
Key questions	3
What is the current situation?.....	3
Are vaccines compulsory for HFEA licensed clinic staff?	3
Should fertility patients be offered the COVID-19 vaccine?	3
What is the current situation regarding treating unvaccinated patients in Scotland?	4
What are The Association of Reproductive and Clinical Scientists (ARCS) and British Fertility Society (BFS) U.K. best practice guidelines for fertility clinics during the COVID-19 pandemic?	4
What if a clinic is unable to continue providing services due to COVID-19?	4
Can clinics continue to offer immunosuppressive treatments?	4
Do patients need to be tested for COVID-19 prior to treatment?	5
Guidance Note 1: Person Responsible	5
What if a PR needs to self-isolate?	5
What if a PR is suddenly unable to fulfil their duties and responsibilities due to COVID-19?	5
What happens if in the interim period between the removal of a PR and the appointment of a new one something occurs at the clinic that requires a PR's urgent decision/action?	5
Guidance Note 2: Staff	5
What guidance should clinic staff follow regarding working from home?	5
What to do if a staff member or patient tests positive for COVID-19.....	6
Guidance Note 3: Counselling and patient support	6
What information can clinics refer their patients to?	6
Can partners attend fertility appointments?.....	7
Guidance Note 4: Information to be provided before consent	7
How should refunds be handled if treatment is cancelled or postponed due to the ongoing COVID-19 pandemic, for example if patients or staff need to self-isolate?	7
Guidance Note 5: Consent to treatment, storage, donation, and disclosure of information	7
Can patients complete and sign their consent forms at home if they cannot come into the clinic? ..	7
Guidance Note 11: Donor recruitment, assessment and screening	8
What guidelines should be referred to for donor screening?	8
Can a donation take place if the gamete donor tests negative for COVID-19?	8
Do patients and donors travelling to England for fertility treatment need to self-isolate?	8
Can a donation take place if the gamete donor has received a COVID-19 vaccine?.....	8
Guidance Note 15: Procuring, processing and transporting gametes and embryos	9

Do people travelling to England to transport gametes or embryos need to self-isolate and produce a negative COVID-19 result?	9
Guidance Note 16: Import and exports	10
Can imports and exports continue as a result of the Coronavirus pandemic?	10
Guidance Note 17: Storage of gametes and embryos	10
Where can I find information on the new Coronavirus Storage Regulations?	10
Under the Coronavirus Storage Regulations, who is eligible to extend their storage to 12 years?	10
Should clinics be contacting all patients who have samples in storage to tell them about the extension to 12 years?	11
Guidance Note 18: Witnessing and assuring patient and donor identification	11
Is it possible to use different measures for witnessing if the number of available clinic staff is limited?	11
Guidance Note 25: Premises, practices and facilities	11
Where can we find information about travel in England?	11
What guidance should be followed regarding COVID-19 laboratory testing?	11
If a patient has had a COVID-19 test performed by another lab and it tested negative, but it is unclear whether this is an accredited test to ISO 15189, is this a valid test or should it be repeated?	11
Does this apply to the new COVID-19 Antibody tests (test to see whether IgG levels to COVID-19 are present) that are currently being evaluated and implemented across UK Medical Laboratories?	12
Can clinics charge for PPE?	12
What procedures should clinics follow if equipment servicing, preventative maintenance or repair cannot be undertaken while the clinic is open, because the clinic's agents are unable to visit due to the pandemic (for instance because of the need to self-isolate)?	12
Guidance Note 32: Obligations and reporting requirements of centres.....	13
What should clinic staff do regarding providing information under EDI timescales?	13
What is the process if a staff member wants to 'whistleblow'?	13
Redundant questions.....	14
What will happen if there is a localised lockdown due to COVID-19?	14
Can I have the vaccine if I work in a private clinic?	14
What is the process for clinics wanting to apply to reopen?	14
What guidance should clinics refer to in preparing the Treatment Commencement Strategy?	15
What is the procedure for completing the self-assessment tool?	15
How will patients know if a centre is open for treatment?	15
What if a PR is furloughed?	15
What should a clinic include in their lone working policy?	15
Are clinic staff 'Key Workers'?	16
Are fertility clinic staff eligible for Coronavirus testing?	16
Is the HFEA making any provisions for extending the 3-month validity for screening blood tests given that cycles have been delayed?	16

Can clinics charge for COVID-19 tests?.....	16
As the Government has relaxed its policy related to screening and has provided a timeframe for the lifting of shielding, can clinics follow Government guidance on shielders or should the BFS/ACRS guidance still be followed?	16

Key questions

What is the current situation?

COVID-19 cases remain high and clinics should continue to follow guidelines around the safe offering of treatment, continue to act carefully, and remain cautious. For information about which recent changes to COVID-19 advice in England see [here](#).

In Scotland: [GOV.SCOT](#).

In Wales: [Read the rules for Wales on GOV.WALES](#).

In Northern Ireland: [Find out more about what you can and cannot do in Northern Ireland on nidirect](#).

Are vaccines compulsory for HFEA licensed clinic staff?

All individuals undertaking [CQC regulated activities in England must](#) be fully vaccinated against COVID-19 no later than 1 April 2022. It is our understanding that the law as it stands today only applies to staff undertaking CQC regulated activities in England.

However, regardless of the legal position, healthcare workers are encouraged to be fully vaccinated against COVID-19. All healthcare leaders are reviewing their staffing in light of this and encouraging patient-facing staff to be fully vaccinated as soon as possible or considering re-deploying non vaccinated staff away from patient-facing activities.

We continue to encourage all patients to get vaccinated in line with recommendations from the BFS, ARCs and RCOG.

Should fertility patients be offered the COVID-19 vaccine?

The Association of Reproductive and Clinical Scientists (ARCS) and [British Fertility Society](#) (BFS) has said that there is absolutely no evidence, and no theoretical reason, that any of the vaccines can affect the fertility of women or men. This includes whether you are trying on your own or having fertility treatment. Recently published [research](#) has supported these claims.

The ARCS and BFS COVID Group issued an [update to the joint guidelines regarding COVID 19 vaccination and fertility](#) on 4 August 2021 and have a set of [FAQs](#) in response to questions that patients have been asking about Covid-19 vaccines and fertility.

Public Health England have also released [guidance on COVID-19 vaccination](#) for women of childbearing age, pregnant or breastfeeding.

More information can be found on the Royal College of Obstetricians and Gynaecologists (RCOG) [website](#).

Vaccination for pregnant women:

The NHS and Royal College of Obstetricians and Gynaecologists (RCOG) has said vaccination against COVID-19 is recommended in pregnancy and is the safest and most effective way of [protecting pregnant women and their babies from coronavirus infection](#).

Pfizer and Moderna vaccines are the preferred vaccines for pregnant women of any age who are coming for their first dose according to the NHS.

More information on this can be found on the [NHS website](#), [RCOG website](#) and [GOV.UK website](#).

What is the current situation regarding treating unvaccinated patients in Scotland?

The Scottish Government has decided that all fertility treatment for COVID-19 unvaccinated women has been deferred where the treatment could result in a pregnancy, with immediate effect (or from whenever is practical). This decision will be reviewed during February 2022 or earlier if appropriate. This includes all fertility treatment except urgent fertility preservation. All patients having treatment deferred will have the deferral time added back on to their treatment journey to ensure that no patient loses out on treatment due to their age. More information can be found [here](#).

What are The Association of Reproductive and Clinical Scientists (ARCS) and British Fertility Society (BFS) U.K. best practice guidelines for fertility clinics during the COVID-19 pandemic?

The latest guidance from the ARCS/BFS COVID working group (August 2021) can be found [here](#).

The guidance sets out five key principles:

- Provision of fertility services must take place in a manner that minimises the chances of spread of COVID-19 infection to patients and fertility clinic staff.
- Centres should ensure a fair and transparent approach to any prioritisation policy.
- Provision of treatment should not result in an undue burden on the NHS.
- Clinical judgement and individualisation of treatment according to the needs of the patient remains at the heart of good clinical care. Patients considering treatment should be fully informed about the effect of the ongoing pandemic on their treatment and give informed consent to having fertility treatment at this time, considering their individual risk profile.
- The fertility sector should adopt sustainable changes in working practices that help to build resilience against any future increases in the spread of COVID-19 in the community.

What if a clinic is unable to continue providing services due to COVID-19?

Local decisions may mean that a centre has to suspend its services. If you decide to suspend services it is important that you communicate clearly with patients, explaining what this means for their treatment, and keep them updated on the situation as it develops. You should also inform your inspector if you decide to suspend services or close your centre.

Can clinics continue to offer immunosuppressive treatments?

There are various different treatments associated with reproductive immunology, which are used to suppress the body's natural immunity, and all of which have risks. These include steroids (e.g. prednisolone), intravenous immunoglobulin (IVIg), TNF- α blocking agents (e.g. adalimumab, infliximab) and intralipid infusions.

Immunosuppressive treatments are not recommended because they are of unproven benefit and carry risks (some of which are serious). They are marked as 'red' on our traffic light system.

[Professional advice](#) from the British Fertility Society, the Association of Reproductive and Clinical Scientists and the Royal College of Obstetricians and Gynaecologists states: While good clinical judgement remains key in the management of individual patients at this time, the use of empirical treatments of uncertain efficacy and safety, including immunosuppressive treatments should be avoided.

Our advice remains that immunosuppressive treatments should not be offered as there is no evidence that these are effective.

Do patients need to be tested for COVID-19 prior to treatment?

Some clinics may choose to include testing for COVID-19 as part of their treatment strategy for all patients having treatment including patients who are asymptomatic. We have not stated clinics must test all patients as it is not a current recommendation made by Government or the Professional Bodies (BFS/ARCS). If you'd like more information on COVID-19 testing, please refer to [the Government guidance](#) which states that anyone with [symptoms](#) can get a polymerase chain reaction (PCR) COVID-19 test, whatever their age. Those with no symptoms can [order rapid lateral flow home test kits online](#).

Guidance Note 1: Person Responsible

What if a PR needs to self-isolate?

If a PR is required to self-isolate because of COVID-19 they would need to decide if they are able to continue working from home during this time or whether they would be unable to continue with their role of PR for the duration of their self-isolation. The PR should contact their inspector to discuss their decision on whether they can continue to work during their self-isolation.

If the PR is unable to fulfil their duties as PR, the Licence Holder must apply for a licence variation with the consent of the person they are proposing to become the new PR. The proposed new PR must also have completed the PREP test. When the PR can resume their duties as PR they would need to be "re-appointed", and the licence variation would need another application to go back to ELP for review.

What if a PR is suddenly unable to fulfil their duties and responsibilities due to COVID-19?

If a PR becomes unwell with symptoms of COVID-19 or is diagnosed with COVID-19 and is suddenly unable to fulfil their duties and responsibilities, they or someone at the clinic must inform their inspector as soon as possible.

Depending on the circumstances, it may be necessary for the Licence Holder to apply for a licence variation to appoint someone else as PR.

What happens if in the interim period between the removal of a PR and the appointment of a new one something occurs at the clinic that requires a PR's urgent decision/action?

If a PR is currently not in their position at the clinic due to COVID-19 and a new PR has not yet been appointed, but an issue or concern has arisen that requires the PR's input, the next in-line manager should contact the centre's inspector or the HFEA incident team with their plan of action. The HFEA would take any necessary regulatory proportionate view on the issue/concern that has arisen.

Guidance Note 2: Staff

What guidance should clinic staff follow regarding working from home?

The Human Fertilisation and Embryology Act 1990 (as amended) ('the Act') does not preclude clinics from allowing certain administrative tasks to be undertaken by clinic staff working from home, where it is

essential for those tasks to be undertaken. However, it is important that clinics ensure appropriate safeguards are put in place to maintain patient confidentiality and that staff who are working from home understand the strict confidentiality provisions which apply both under section 33A of the Act and under Data Protection legislation. Before permitting clinic staff to work from home, clinics should develop appropriate home working and security policies in consultation with IT staff, data protection advisors and/or legal advisors. Consideration should be given to matters including:

1. access to patient data or records over secure networks and ideally access should only be via a laptop or computer provided by the clinic which is password protected and encrypted;
2. staff understanding of the confidentiality requirements of section 33A of the Act and a reminder that disclosing patient data in breach of section 33A is a criminal offence;
3. Data Protection requirements that might apply when accessing patient data or patient records from home;
4. practical steps that staff should take when working from home to avoid inadvertent breaches of patient confidentiality or unlawful disclosure of patient records. For example, preventing staff from being able to print out any patient records or patient data at home;
5. measures that can be taken to ensure that other people at the staff member's home cannot see or access any patient data or patient records on the staff member's work computer or laptop;
6. preventing staff members from taking hard copy patient records home;
7. reminding staff members to ensure that personal or private medical matters relating to a patient are not discussed within earshot of their family or friends, ideally (if possible) one would want staff to have a private space in their home where they can work away from family etc.;
8. the need to develop a "Data Protection and Patient Confidentiality Policy for home workers" which sets out what is expected of staff, what they can and cannot do etc. and what they need to do if there is unlawful access to patient data or an inadvertent data protection breach;
9. the need to comply with the usual information security arrangements that are in place in clinics whilst in the home environment. For example, clear desks, locking screens when away from the desk and storing laptops securely when not in use;
10. what staff members are expected to do if there is a data breach or if they suspect that the security of their network or device may have been compromised by malware or unauthorised hacking or other IT security event;
11. the importance of ensuring that all email communication is addressed to the correct recipient(s) before it is sent, and that appropriate security measures are put in place if staff need to send sensitive personal data/patient records electronically. For example, two factor authentications (password sent via telephone or in separate email).

What to do if a staff member or patient tests positive for COVID-19.

If any staff member or patient tests positive for COVID-19, this must be reported to the HFEA via the HFEA incident reporting system and investigated appropriately. Centres should ensure cases or suspected cases are reported appropriately to facilitate the [Track and Trace](#) programme. This will also ensure early identification of an outbreak within a centre.

Guidance Note 3: Counselling and patient support

What information can clinics refer their patients to?

There are websites that can provide support and/or information to patients at this time of uncertainty including:

- [HFEA Coronavirus \(COVID-19\) guidance for patients](#)

- [Fertility Network UK](#)
- [British Infertility Counselling Association](#)
- [Fertility Friends](#)
- [Coronavirus \(COVID-19\): what you need to do \(Government advice\)](#)
- [NHS advice](#)

Centres should ensure patients have access to information regarding the effects of coronavirus infection during pregnancy. Each clinic should state if they are able to offer telephone or online counselling to their patients.

Can partners attend fertility appointments?

Centres may implement their own policies and different decisions may be made at a clinic level based on local circumstances about any measures they have in place, for example, when allowing partners to attend appointments.

The NHS has introduced a [framework](#) to allow partners to be able to attend maternity services in England, including fertility treatments. They recommend using a stepwise approach, following a documented risk assessment, so you can make any necessary changes before relaxing any current approaches you may have in place. Your policies on permitting access to partners should be regularly reviewed, be tailored to your local context and take account of current risks and government policy.

The [updated guidance \(4 August 2021\)](#) from The Association of Reproductive and Clinical Scientists (ARCS) and British Fertility Society (BFS) says that centres should minimise the number of accompanying persons, and should carry out risk assessments to ensure the attendance of a partner at appointments is safe.

[NHS guidance](#) says that further to a risk assessment, women should have access to support from a person of her choosing at all stages of her maternity journey and that all trusts should facilitate this as quickly as possible.

Guidance Note 4: Information to be provided before consent

How should refunds be handled if treatment is cancelled or postponed due to the ongoing COVID-19 pandemic, for example if patients or staff need to self-isolate?

Before a patient begins treatment, consumer protection law states that clinics should explain in writing the circumstances in which treatment may be cancelled or postponed, and explain a patient's rights and obligations should treatment be cancelled or postponed e.g. whether they are entitled to a refund of any prepayments.

The Competition and Markets Authority (CMA) has published [guidance](#) which sets out its views in relation to cancellations and refunds within the context of the COVID-19 pandemic. This explains the circumstances in which the CMA would expect a consumer to be offered a full refund where no services are provided as a result of COVID-19. This guidance also explains that in circumstances where a patient has already received some services they have paid for in advance, that they would be entitled to a refund of the services not already provided.

Guidance Note 5: Consent to treatment, storage, donation, and disclosure of information

Can patients complete and sign their consent forms at home if they cannot

come into the clinic?

In the Code of Practice, we provide guidance on obtaining consent in Guidance Note 5.11 and Consent should be given at the clinic (with both parties present if a couple is being treated) or a documented process should be in place to ensure that consent forms signed outside the clinic are signed by the correct person, have been correctly completed and the consent is valid. Therefore, if clinics have a documented process in place to ensure the identity of their patients, and to ensure the forms are completed correctly, consents can be signed outside of the clinic. Detailed guidance on introducing electronic methods of taking consent can be found in Guidance Note 5.31- 5.38.

Guidance Note 11: Donor recruitment, assessment and screening

What guidelines should be referred to for donor screening?

The Advisory Committee on the Safety of Blood, Tissues and Organs (SaBTO) updated their guidelines on [Microbiological Safety Guidelines](#). Please refer to Annex 1: Organs, tissue and cells used in transplantation and SARSCoV-2 (page 61), which contains guidance for gamete donors. In addition to this there is signposting to [NHSBT guidance for tissues and organs](#).

The following applies:

Symptomatic gamete donor – The donor should be deferred if, at the time of donation or in the preceding 7 days, the donor has a new continuous cough or a temperature $>37.8^{\circ}\text{C}$. The donor can donate after 28 days from full recovery.

Asymptomatic gamete donor – If the donor is asymptomatic, government advice about those who should self-isolate should be considered including the implications this may have on the donation process.

JPAC (Joint United Kingdom (UK) Blood Transfusion and Tissue Transplantation Services Professional Advisory Committee) issued a Change Notification regarding live tissue, which states also that person with confirmed symptomatic Coronavirus infection must not donate if less than 28 days since resolution of symptoms. All latest guidance can be found on the [JPAC website](#).

Can a donation take place if the gamete donor tests negative for COVID-19?

If an apparent symptomatic donor gamete is tested for COVID-19 and tests negative, the current SaBTO advice should still be followed as expert consultation would be needed on the diagnostic reliability of tests at different stages of infection before making any change.

Do patients and donors travelling to England for fertility treatment need to self-isolate?

What visitors to England must do depends on whether individuals qualify as fully vaccinated and on whether they have been in a red list country in the 10 days prior to arriving in England.

More information can be found here: [Travel to England from another country during coronavirus \(COVID-19\)](#).

Can a donation take place if the gamete donor has received a COVID-19 vaccine?

If donors have received a COVID-19 vaccine in the UK vaccination, they must not donate if less than seven days have passed after the most recent vaccination was given. If the donor feels unwell after the vaccination, they must not donate for 7 days after the resolution of symptoms.

If donors have received a COVID-19 vaccine outside the UK vaccination programme, including participants in clinical trials or donors vaccinated outside the UK, clinics should refer to a Designated Clinical Support Officer for individual risk assessment. Additional information can be found [here](#) in change notification no 05.

Guidance Note 15: Procuring, processing and transporting gametes and embryos

Do people travelling to England to transport gametes or embryos need to self-isolate and produce a negative COVID-19 result?

The rules determining what people travelling to England to transport gametes or embryos must do before entering England depends on whether they are vaccinated or unvaccinated, and on whether they have been in a red list country in the 10 days before arriving in England.

Non-red list countries

Before you travel to the UK you need to [complete a passenger locator form](#). You do not need to take a COVID-19 test before you travel to England. You do not need to book COVID-19 tests to take after arrival in England.

If you are fully vaccinated, you do not need to quarantine but you must take a lateral flow device (LFD) test on or before day 2 of your visit. If you are travelling internationally on a daily basis, you need to take an LFD test every 3 days.

If you are not fully vaccinated and you will be in England for longer than 2 days, you need to take an LFD test on or before days 2, 5 and 8 after your arrival. You must [quarantine in the place you are staying for 10 days](#). You can leave your accommodation when conducting a specific exempt activity or travelling directly to and from your accommodation and a location for the purposes of this work. You must quarantine at all other times during your stay.

If you are travelling every day, you must take a lateral flow test at least once every 3 days.

Red list countries:

Applies if you have been in a [red list country](#) in the 10 days before you arrive in England - you do not need to quarantine in a government approved quarantine hotel.

You must [quarantine in the place you are staying](#). You can leave your accommodation when conducting the specific exempt activity or travelling directly to and from your accommodation and a location for the purposes of this work. You must quarantine at all other times during your stay. When you are in a red list country you should take additional precautions to minimise contact with others.

If you are in England for longer than 2 days, you need to take a mandatory workplace, community or home [LFD test](#) on or before days 2, 5 and 8 after arrival.

If you are travelling every day, you must take a lateral flow test at least once every 3 days.

Evidence: will need to show:

- staff pass or photo identification
- confirmation from a healthcare provider explaining:
 - the address of where the material was or is to be collected from, and the date on which the material was or is to be collected
 - the address of where the material is to be transported to, the name of the healthcare provider that is to receive the material, and the date on which the material is to be delivered to the healthcare provider

- what material you are or will be carrying
- why you need to carry it

Exemption from COVID-19 testing before entering England

You do not need to present a negative coronavirus result test prior to departure.

You will need to show:

- your staff pass or photo identification
- confirmation from a healthcare provider explaining:
 - the address of where the material was or is to be collected from, and the date on which the material was or is to be collected
 - the address of where the material is to be transported to, the name of the healthcare provider that is to receive the material, and the date on which the material is to be delivered to the healthcare provider
- what material you are or will be carrying
- why you need to carry it

For more information on the rules in England, see [here](#).

Scotland: <https://www.gov.scot/publications/coronavirus-covid-19-international-travel-quarantine/>

Northern Ireland: <https://www.nidirect.gov.uk/articles/coronavirus-covid-19-international-travel-advice>

Wales: <https://gov.wales/foreign-travel-and-returning-home>

Guidance Note 16: Import and exports

Can imports and exports continue as a result of the Coronavirus pandemic?

HFEA requirements and guidance pertaining to gamete and embryo imports have not changed during the COVID-19 emergency. While treatment was suspended during the emergency by General Direction 0014, a suspension of imports and exports was not required by the HFEA and therefore can continue as normal.

Guidance Note 17: Storage of gametes and embryos

Where can I find information on the new Coronavirus Storage Regulations?

On 1 July 2020, the Government changed the law to extend the current 10-year storage limit for eggs, sperm and embryos by 2 years so that those undergoing fertility treatment during the coronavirus outbreak have more time to continue treatment. The Regulations can be found here: [The Human Fertilisation and Embryology \(Statutory Storage Period for Embryos and Gametes\) \(Coronavirus\) Regulations 2020](#).

We issued a Chair's letter and FAQs for clinics, which can be found [here](#) and a new consent form (the CVS form) which can be found [here](#).

Under the Coronavirus Storage Regulations, who is eligible to extend their storage to 12 years?

Patients whose gametes or embryos are in storage at a licensed treatment, storage or research clinic are able to extend their storage for 12 years provided:

- their gametes or embryos were in storage on 1 July 2020, the date the Regulations came into force, and
- they consent in writing to their gametes or embryos being stored for at least 12 years. This consent can be provided before, on or after 1 July 2020.

Should clinics be contacting all patients who have samples in storage to tell them about the extension to 12 years?

It is a local decision as to when clinics decide to contact their patients to tell them about the possibility of extension; they may choose to do this now or when their patients are nearing the end of their initial storage period. Sometimes it can be challenging to contact patients, and this should be taken into account when centres decide when to contact patients about this complex consenting situation, so as to ensure effective storage consent is continuously documented for all stored samples. Patients who may have been affected by clinic closures should be prioritised.

Guidance Note 18: Witnessing and assuring patient and donor identification

Is it possible to use different measures for witnessing if the number of available clinic staff is limited?

Licence conditions and the Code of Practice require a witnessing protocol to be in place, and a robust risk assessment to be carried out if the protocol is changed. Due to current Government recommendations for workers that are able to do so to work from home, the use of live video conferencing as an alternative witnessing protocol is acceptable. Before any new protocol is used, it should be risk assessed to allow any potential problems to be identified, such as who is in the room with the remote witness, and whether or not the video conferencing platform is encrypted.

Guidance Note 25: Premises, practices and facilities

Where can we find information about travel in England?

What visitors to England must do depends on whether individuals qualify as fully vaccinated and on whether they have been in a red list country in the 10 days prior to arriving in England.

More information can be found here: [Travel to England from another country during coronavirus \(COVID-19\)](#).

What guidance should be followed regarding COVID-19 laboratory testing?

- NHS: [Guidance and standard operating procedure COVID-19 virus testing in NHS laboratories](#)
- Department of Health and Social Care: [Guidance for organisations seeking to support the COVID 19 Testing](#)
- Department of Health and Social Care: [Coronavirus \(COVID-19\): Scaling up our testing programmes](#)

If a patient has had a COVID-19 test performed by another lab and it tested negative, but it is unclear whether this is an accredited test to ISO 15189, is this a valid test or should it be repeated?

All tests that have been performed via the government approved route are valid tests. There should be no need to repeat the test if the patient tested negative unless there was concern that there had been recent exposure since the last test. For more information, please see [Guidance and standard](#)

[operating procedure COVID-19 virus testing in NHS laboratories](#) for more information.

The government, NHSE, and DHSC issued guidelines regarding the scope of accreditation for this test and in accordance with Public Health England (PHE) have devised strict validation processes to ensure the test is accurate and reliable.

Does this apply to the new COVID-19 Antibody tests (test to see whether IgG levels to COVID-19 are present) that are currently being evaluated and implemented across UK Medical Laboratories?

Yes, this guidance is also applicable to the COVID-19 Antibody test. Although not all laboratories have a supply of these reagents as their supply is being nationally coordinated by NHSE and DHSC.

It is important to follow all government NHS/DHSC/PHE guidance regarding Covid-19 testing to ensure the most recent guidelines regarding testing are being adhered to. Please refer to relevant government guidelines for the current validated tests available.

Can clinics charge for PPE?

Covering costs is acceptable but any price should be fair.

What procedures should clinics follow if equipment servicing, preventative maintenance or repair cannot be undertaken while the clinic is open, because the clinic's agents are unable to visit due to the pandemic (for instance because of the need to self-isolate)?

Some servicing and repair services may be harder to access when the pandemic is more active. If your usual servicing and repair agents cannot or will not visit your clinic, in the first instance you must try to use an appropriate alternative agent, especially for critical equipment. Efforts to find an alternative agent should be documented.

If no alternative agents are available:

For equipment beyond its service interval: Clinics may continue using such equipment provided:

- The clinic monitors the equipment continually, or at an appropriate frequency, and there are no significant or unexplained non-conformances in performance, OR an in-house re-validation of the equipment (as a minimum an EQR) is performed before use in treatment, without significant or unexplained non-conformances in performance. The review of monitoring/validation of the equipment must be completed by a staff member who has the technical knowledge to do so and must be fully documented.
- The clinic must arrange for a service visit to be completed by an appropriate agent, within three months of such an agent becoming available. If this is not possible because of demand on the agent's services, you should document all correspondence and contacts on the matter and should arrange a servicing visit at the earliest opportunity.

For equipment requiring a preventative maintenance visit: Monitoring may allow equipment to be used past a scheduled preventive maintenance visit in some cases but manufacturers are best placed to assess whether a piece of equipment can be safely used. Where the recommended preventative maintenance period on a clinic's equipment has expired and a visit cannot be arranged, clinics may continue using such equipment provided:

- The clinic monitors the equipment continually, or at an appropriate frequency, and there are no significant or unexplained non-conformances in performance, OR an in-house re-validation of the equipment (as a minimum an EQR) is performed before use in treatment, without significant or

unexplained non-conformances in performance. The review of monitoring/validation of the equipment must be completed by a staff member who has the technical knowledge to do so and must be fully documented.

- The clinic has written confirmation from the equipment manufacturer that operation of the equipment beyond the recommended preventative maintenance period is highly unlikely, for a defined period, to undermine equipment performance. Use of the equipment in treatment should cease before the end of that defined period.
- The clinic assesses the risks of continuing to use equipment requiring a preventative maintenance visit and, where possible, takes actions to mitigate any risks identified.
- The clinic must arrange for a preventative maintenance visit to be completed by an appropriate agent, within three months of such an agent becoming available. If this is not possible because of demand on the agent's services, you should document all correspondence and contacts on the matter, and should arrange a preventative maintenance visit at the earliest opportunity.

For equipment which develops a fault: Failure to repair equipment may lead to out of specification performance or failure, but it is also possible that a fault in a non-essential system may not undermine performance. The effect of a fault on equipment performance will depend on multiple factors, best considered by the manufacturer. Thus where equipment develops a fault which repair engineers cannot visit to address, the equipment can continue to be used only if:

- The clinic monitors the equipment continually, or at an appropriate frequency, and there are no significant or unexplained non-conformances in performance, OR an in-house re-validation of the equipment (as a minimum an EQR) is performed before use in treatment, without significant or unexplained non-conformances in performance. The review of monitoring/validation of the equipment must be completed by a staff member who has the technical knowledge to do so and must be fully documented.
- The clinic has written confirmation from the equipment manufacturer that operation of the equipment with the fault, is highly unlikely, for a defined period, to undermine equipment performance. Use of the equipment in treatment should cease before the end of that defined period.
- The clinic assesses the risks of continuing to use equipment with identified faults and, where possible, takes actions to mitigate any risks identified.
- The clinic must arrange for a repair visit to be completed by an appropriate agent, within three months of such an agent becoming available. If this is not possible because of demand on the agent's services, you should document all correspondence and contacts on the matter and should arrange a repair visit at the earliest opportunity.

Guidance Note 32: Obligations and reporting requirements of centres

What should clinic staff do regarding providing information under EDI timescales?

We will not necessarily take regulatory action against any clinics that fail to comply with the requirements regarding the provision of information on EDI forms where their failure to do so is directly attributable to COVID-19 or circumstances arising as a consequence of it. The clinic must assure us that the data is being collected, to be collated and reported once the clinic is able to resume its normal functions.

Clinics must not post any forms required for providing information to us until further notice because HFEA staff are working from home as of December 2021. For further guidance about what to do instead of sending urgent postal correspondence, please send an email to enquiriesteam@hfea.gov.uk.

What is the process if a staff member wants to 'whistleblow'?

Clinic staff are aware that they are able to use our incident email (Incident.Reporting@HFEA.GOV.UK) to whistle blow if they have any concerns about working practices within their clinics during the Covid-19 emergency.

Redundant questions

What will happen if there is a localised lockdown due to COVID-19?

If there is a localised lockdown, it will be the responsibility of the PR to assess the risk locally, to decide whether treatment can continue to take place for patients. This decision should be informed by knowledge of local coronavirus infection rates and local measures, if any. This would involve looking at the risks to patients and would need to take into account factors such as the local 'r' rate and any restrictions in place locally. For a centre to be approved to reopen the revised General Direction 0014v2 required centres to have a written Covid-19 Treatment Commencement Strategy. This means clinics will have already put measures in place to ensure they comply with specified guidance on safe and effective working practices during the pandemic. A PR will need to assess whether these are sufficient to be able to protect patients in light of a new lockdown. The HFEA will continue to monitor national and devolved government advice as well as that of the professional bodies and will react in an appropriate manner, should the situation change at any time.

It should be recognised that patients are likely to be anxious about coronavirus infection and its potential effects on pregnancy. Patients should be made aware that the present experience is limited and does not indicate that the severity of infection is any worse in pregnancy. At this stage, there is no evidence of an increased risk of fetal anomalies or adverse pregnancy complications. Nonetheless, patients should be carefully counselled, taking into account their individual clinical situation and risk profile, and the likely persistence of the virus in the local community in the medium term. This counselling and the patient's decision whether or not to proceed with fertility treatment should be documented in the medical record.

Can I have the vaccine if I work in a private clinic?

The [COVID-19 Greenbook](#) (updated 1 July 2021) states that priority group 2 includes frontline healthcare staff, this includes staff who have frequent face-to-face clinical contact with patients and who are directly involved in patient care in either secondary or primary care. This includes doctors, midwives and nurses. It should also include those working in independent, voluntary and non-standard healthcare settings. This group also includes non-clinical staff in secondary or primary care settings, including non-clinical ancillary staff who may have social contact with patients but are not directly involved in patient care. This group includes receptionists, ward clerks, porters and cleaners. Also included are hospital-based laboratory staff who frequently handle SARS-CoV-2 or collect or handle potentially infected specimens including blood specimens should be eligible as they may also have social contact with patients. This may also include cleaners, porters, secretaries and receptionists in laboratories. Staff working in non-hospital-based laboratories and those academic or commercial research laboratories who handle clinical specimens or potentially infected samples will be able to use effective protective equipment in their work and should be at low risk of exposure.

What is the process for clinics wanting to apply to reopen?

The HFEA issued revised [GD0014 \(version 2\)](#) on 11 May 2020 which enables any licensed centre to apply to resume treatment services provided it can satisfy certain safety requirements. Clinics are required to have a COVID-19 Treatment Commencement Strategy in order to be able to apply to reopen.

The Strategy records the measures the clinic will be taking to comply with current professional body guidance on safe and effective treatment; records the risk assessments undertaken; the practical and logistical arrangements that will be put in place at the clinic and records all new or

revised standard operating procedures (SOPs) or protocols.

The revised GD0014 also requires the clinic to complete the HFEA COVID-19 Treatment Commencement Self-assessment which has been developed to measure the robustness of the clinic's Treatment Commencement Strategy and assess the clinic's compliance with guidance from the UK and devolved governments, professional bodies such as [BFS/ARCS](#), ESHRE, as well as Standard Licence Conditions and guidance in the Code of Practice. Before a clinic can commence treatment, it must have received approval of its self-assessment from its HFEA inspector. Any clinic that commences treatment before having received this approval will be in breach of GD0014 and may face regulatory sanction.

When you are ready to apply to re-open, the clinic's HFEA inspector should be contacted.

What guidance should clinics refer to in preparing the Treatment Commencement Strategy?

The following guidance should be referred to:

- [British Fertility Society – COVID-19](#)
- [GOV.UK – COVID-19: infection prevention and control \(IPC\)](#)
- [GOV.UK – Getting tested](#)
- [ESHRE – COVID-19](#)

This list is not exhaustive, and PRs should incorporate the sections of guidance relevant to local practice. You should keep your Treatment Commencement Strategy under regular review to ensure that it reflects the latest guidance.

What is the procedure for completing the self-assessment tool?

The tool is available from the clinic's HFEA inspector. It is a questionnaire containing 50 questions. The completed questionnaire should be sent to the clinic's inspector who will review the responses within five working days with a view to the proposal to commence treatment services. We aim to complete our approval process within 5 working days of receiving the completed clinic self-assessment.

How will patients know if a centre is open for treatment?

As centres receive approval to reopen, we will publish this information in a list on our website to make patients aware. This list can be found [here](#).

What if a PR is furloughed?

PRs who have been furloughed due to COVID-19 may remain PR provided they can continue to fulfil their statutory duties as PR during furlough and do not contravene the government's rules around furlough. If this is not possible, it may be necessary for the licence to be varied to appoint someone else as PR.

There are no statutory restrictions on a person becoming PR of more than one clinic. If it is necessary for someone else to be appointed as PR, the Licence Holder will need to apply for a licence variation with the consent of the person they are proposing to become the new PR.

The proposed new PR must also have completed the PR Entry Programme (PREP) test. The HFEA will expedite such applications so they can be dealt with as quickly as possible. When the HFEA allows clinics to resume treatment, clinics will need to consider whether the former PR wishes to resume his/her role as PR and if so, a further application will need to be made to the HFEA.

What should a clinic include in their lone working policy?

Each clinic should have updated their lone working policy to now include any aspects specific to lone working during the COVID-19 pandemic. This should include but is not limited to;

- a procedure for staff to follow if they are working on their own e.g. a telephone number of someone they have to notify when they arrive and when they leave
- an updated safety procedure if an emergency or accident occurs whilst someone is working alone, this should include emergency contacts
- safety of staff who need to perform certain duties e.g. maintaining the liquid nitrogen dewars
- use of PPE by all staff
- planning to contact the providers (of liquid nitrogen etc) to discuss any supply chain issues, when they should be placing their order and ensure that the provider knows that the clinic would be considered a critical service

Are clinic staff 'Key Workers'?

Public Health England have confirmed that any healthcare staff engaged in the completion of current fertility treatment cycles are considered Key workers.

Are fertility clinic staff eligible for Coronavirus testing?

According to the latest [guidance on testing](#), clinic staff are covered under testing where they are showing symptoms or someone they live with is showing symptoms. Anyone providing a medical service is classified as a key worker, including:

- doctors, nurses, midwives, paramedics, social workers, care workers, and other frontline health and social care staff including volunteers
- support and specialist staff required to maintain the UK's health and social care sector
- those working as part of the health and social care supply chain, including producers and distributors of medicines and medical and personal protective equipment.

Is the HFEA making any provisions for extending the 3-month validity for screening blood tests given that cycles have been delayed?

The screening requirements for patients undertaking IVF are set out in the Act (and most originate within the EUTCD).

Because of the requirement to cease treatment imposed by General Direction 0014, clinics may not be able to undertake the necessary screening within the required timeframes. This may result in the validity of certain screening tests being compromised as the gap between screening blood sampling and treatment is extended beyond three months for those patients planning a first treatment cycle, or two years for those attending for further treatment. Whilst it remains a legal requirement for clinics to satisfy all screening requirements, the HFEA will adopt a proportionate approach to any regulatory action where the time between screening blood sampling and treatment has extended beyond what is permitted because of the requirement to cease treatment by virtue of General Direction 0014, or other circumstances arising as a consequence of COVID-19 or that are directly attributable to COVID-19, provided a risk assessment indicates that the safety of patients and storage systems has not been compromised and the patient is fully informed of the non-compliance.

Can clinics charge for COVID-19 tests?

The current guidance is that, with reference to Government and professional body guidance, centres should consider implementing a testing policy as reliable serological tests become more widely available. **If a patient has symptoms of COVID-19**, you should inform them that they can access a test on the NHS website.

As the Government has relaxed its policy related to screening and has provided a timeframe for the lifting of shielding, can clinics follow

Government guidance on shielders or should the BFS/ACRS guidance still be followed?

The [BFS/ARCS guidance](#) states that 'particular caution should apply to patients with underlying medical problems whose co-morbidity places them at a higher risk of complications in the event of contracting coronavirus infection. This includes patients with obesity, hypertension, diabetes and those receiving immunosuppressive medication. It may be appropriate for such patients to delay conception until epidemiological evidence shows a sustained reduction in the community spread of the infection'. This echoed the government advice that people who are vulnerable from Covid-19 should shield.

On 22 June the government set out a series of steps for further relaxing shielding guidance which will come into effect on 6 July and 1 August. From the 1st August shielding will be paused. The guidance does state that caution and strict social distancing is required, and the advice is still to stay at home where possible. We would therefore expect clinics to assess patients classed as vulnerable on a case by case basis and risk assess how and if treatment can be undertaken safely.